arXiv:1605.03516v1 [math.FA] 11 May 2016

SOME INEQUALITIES FOR THE MATRIX HERON MEAN

DINH TRUNG HOA

Abstract. Let A, B be positive definite matrices, $p = 1, 2$ and $r \geq 0$. It is shown that

$$||A + B + r(A\sharp_t B + A\sharp_{1-t} B)||_p \leq ||A + B + r(A^t B^{1-t} + A^{1-t} B^t)||_p.$$

We also prove that for positive definite matrices A and B

$$\det(P_t(A, B)) \leq \det(Q_t(A, B)),$$

where $Q_t(A, B) = \left(\frac{A^t + B^t}{2}\right)^{1/t}$ and $P_t(A, B)$ is the t-power mean of A and B. As a consequence, we obtain the determinant inequality for the matrix Heron mean: for any positive definite matrices A and B,

$$\det(A + B + 2(A\sharp B)) \leq \det(A + B + A^{1/2}B^{1/2} + A^{1/2}B^{1/2})).$$

These results complement those obtained by Bhatia, Lim and Yamazaki (LAA, **501** (2016) 112-122).

1. Introduction

Let M_n be the space of $n \times n$ complex matrices and M_n^+ the positive part of M_n. Denote by I the identity element of M_n. For self-adjoint matrices $A, B \in M_n$ the notation $A \leq B$ means that $B - A \in M_n^+$. For a real-valued function f of a real variable and a self-adjoint matrix $A \in M_n$, the value $f(A)$ is understood by means of the functional calculus.

For $0 \leq t \leq 1$ the t-geometric mean of A and B is defined as

$$A\sharp_t B = A^{1/2}(A^{-1/2}BA^{-1/2})^t A^{1/2}.$$

The geometric mean $A\sharp B := A\sharp_{1/2} B$ is the midpoint of the unique geodesic $A\sharp_t B$ connecting two points A and B in the Riemannian manifold of positive matrices.

Recently, Bhatia et al. [1] proved that for any positive definite matrices A and B and for $p = 1, 2$,

$$||A + B + 2rA\sharp B||_p \leq ||A + B + r(A^{1/2}B^{1/2} + B^{1/2}A^{1/2})||_p. \tag{1}$$

When $r = 1$, inequality holds for $p = \infty$.

For the case $p = 2$ the proof of (1) is based on the following fact: for any positive definite A and B,

$$\lambda(A^{1/2}(A\sharp B)A^{1/2}) \prec_{\log} \lambda(A^{3/4}B^{1/2}A^{3/4}), \tag{2}$$

2000 *Mathematics Subject Classification.* 46L30, 15A45, 15B57.

Key words and phrases. operator (r, s)-convex functions, operator Jensen type inequality, operator Hansen-Pedersen type inequality, operator Popoviciu inequality.

where the notation λ is used for the n-tuple of eigenvalues of a matrix A in decent order and $\lambda(A) \prec_{\log} \lambda(B)$ means that

$$\prod_{j=1}^{k} \lambda_i(A) \leq \prod_{j=1}^{k} \lambda_i(B), \quad 1 \leq k \leq n$$

and inequality holds when $k = n$.

For $p = \infty$ inequality (1) was proved by using a result of Lim and Yamazaki [3, Theorem 4.1]

$$||P_t(A_1, A_2, \cdots, A_n)||_\infty \leq ||Q_t(A_1, A_2, \cdots, A_n)||_\infty, \tag{3}$$

where *the power mean* $P_t(A_1, A_2, \cdots, A_n)$ of $A_1, A_2, \cdots, A_n$ [2] and is the unique solution of the matrix equation

$$X = \frac{1}{m} \sum_{i=1}^{m} X \sharp_t A_i$$

and

$$Q_t(A_1, A_2, \cdots, A_n) = \big(\frac{1}{m} \sum_{i=1}^{m} A_i^t\big)^{1/t}.$$

For $m = 2$ Lim and Páflia [2, Remark 3.10] show that

$$P_t(A, B) = A\sharp_{1/t}\big(\frac{1}{2}(A + A\sharp_t B)\big) = A^{1/2}\big(\frac{I + (A^{-1/2}BA^{-1/2})^t}{2}\big)^{1/t} A^{1/2}.$$

Hopefully, for $t = 1/2$,

$$P_{1/2}(A, B) = \frac{1}{4}(A + B + A\sharp B), \quad Q_{1/2}(A, B) = \frac{1}{4}(A + B + A^{1/2}B^{1/2} + B^{1/2}A^{1/2}).$$

And so, the inequality (1) for $p = \infty$ is obtained from (3) choosing $m = 2$ and $t = 1/2$.

Recall that the family of Heron mean [6] for nonnegative number a, b is defined as

$$H_t(a, b) = (1 - t)(\frac{a + b}{2}) + t\sqrt{ab}, \quad 0 \leq t \leq 1. \tag{4}$$

The Kubo-Ando extension of this to matrices is

$$(1 - t)\frac{A + B}{2} + t(A\sharp B)$$

that connects the arithmetic mean and the geometric mean, and a naive extension is

$$(1 - t)\frac{A + B}{2} + t\frac{A^{1/2}B^{1/2} + B^{1/2}A^{1/2}}{2}$$

that connects the arithmetic mean and the midpoint of the Heinz mean $\frac{A^sB^{1-s} + B^sA^{1-s}}{2}$. So, inequality (1) is a special case of the following

$$||(1 - t)\frac{A + B}{2} + t(A\sharp B)||_p \leq ||(1 - t)\frac{A + B}{2} + t\frac{A^{1/2}B^{1/2} + B^{1/2}A^{1/2}}{2}||_p$$

with $t = 1/2$.

Notice that another naive extension of the Heron mean for positive definite matrices A and B is defined as

$$(1 - t)\frac{A + B}{2} + t\frac{A^tB^{1-t} + A^{1-t}B^t}{2}.$$

In this paper, we extend the inequality (2) to t-geometric means. More precisely, we prove that for any positive definite matrices A, B and for any $t \in [0,1]$

$$\lambda(A^{1/2}(A\sharp_t B)A^{1/2}) \prec_{\log} \lambda(A^{1-t}B^tA^{1-t}).$$

Using this extension, we prove the following result:

Theorem 1.1. *Let A, B be positive definite matrices and $p = 1, 2$ and $r \geq 0$. Then*

$$||A + B + r(A\sharp_t B + A\sharp_{1-t} B)||_p \leq ||A + B + r(A^tB^{1-t} + A^{1-t}B^t)||_p. \tag{5}$$

Also, using the approach in [4] we show that for positive definite matrices A, B and for any z in the strips $S_{1/4} = \{z \in \mathbb{C} : \mathrm{Re}(z) \in [1/4, 3/4]\}$,

$$|\operatorname{Tr}(A^{1/2}B^zA^{1/2}B^{1-z})| \leq \operatorname{Tr}(AB).$$

2. Inequalities

Proposition 2.1. *Let A, B be positive definite matrices. Then for any $t \in [0,1]$*

$$\lambda(A^{1/2}(A\sharp_t B)A^{1/2}) \prec_{\log} \lambda(A^{1-t/2}B^tA^{1-t/2}). \tag{6}$$

Proof. Firstly, let's prove

$$\lambda_1(A^{1/2}(A\sharp_t B)A^{1/2}) \leq \lambda_1(A^{1-t/2}B^tA^{1-t/2}). \tag{7}$$

This inequality is equivalent to the statement

$$A^{1-t/2}B^tA^{1-t/2} \leq I \quad \Longrightarrow \quad A^{1/2}(A\sharp_t B)A^{1/2} \leq I,$$

which in turn is equivalent to

$$B^t \leq A^{t-2} \quad \Longrightarrow \quad (A^{-1/2}BA^{-1/2})^t \leq A^{-2}. \tag{8}$$

That can be proved by using the Furuta inequality which states that if $0 \leq Y \leq X$, then for all $p \geq 1$ and $r \geq 0$ we have

$$(X^rY^pX^r)^{1/2} \leq (X^{p+2r})^{1/p}. \tag{9}$$

Let apply (9) to $X = A^{t-2}$, $Y = B^t$, $p = \frac{1}{t}$ and $r = -\frac{1}{2(t-2)}$, we get (8), and hence (7).

Denote by $C_k(X)$ the k-th compound of $X \in M_n$, $k = 1, \ldots, n$. Note that for any positive definite matrices X, Y,

$$\begin{aligned} C_k(A^{1/2}(A\sharp_t B)A^{1/2}) &= C_k(A(A^{-1/2}BA^{-1/2})^tA) \\ &= C_k(A)C_k((A^{-1/2}BA^{-1/2})^t)C_k(A) \\ &= C_k^{1/2}(A)(C_k^{1/2}(A)(C_k^{-1/2}(A)C_k(B)C_k^{-1/2}(A))^tC_k^{1/2}(A))C_k^{1/2}(A) \\ &= C_k^{1/2}(A)(C_k(A)\sharp_t C_k(B))C_k^{1/2}(A). \end{aligned} \tag{10}$$

In the other hand,

$$\lambda_1(C_k(A^{1/2}(A\sharp_t B)A^{1/2})) = \prod_{i=1}^{k}\lambda_i(A^{1/2}(A\sharp_t B)A^{1/2}), \quad k = 1, \ldots, n-1, . \tag{11}$$

On account of (7) and (11) for $1 \leq k \leq n$ we have

$$\begin{aligned}\prod_{i=1}^{k}\lambda_i(A^{1/2}(A\sharp_t B)A^{1/2}) &= \lambda_1(C_k(A^{1/2}(A\sharp_t B)A^{1/2})) \\ &\leq \lambda_1(C_k(A)^{1-t}C_k(B)^t C_k(A)^{1-t}) \\ &= \prod_{i=1}^{k}\lambda_i(A^{1-t}B^t A^{1-t}).\end{aligned}$$

The equality holds for $k = n$, since $\det(A^{1/2}(A\sharp_t B)A^{1/2)} = \det(A^{1-t}B^t A^{1-t})$.

Thus, we have proved (6). □

The following special case of Proposition will be used in the proof of the main result.

Corollary 2.2. *For any positive definite matrices A and B,*

$$\mathrm{Tr}(A(A\sharp_t B)) \leq \mathrm{Tr}(A^{2-t}B^t), \quad t \in [0, 1].$$

In order to prove the next result, let's recall the generalized Hølder inequality for trace [5, Theorem 2.8]: let $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$ for $p, q, r \geq 1$ and X, Y, Z be matrices in M_n, then

$$\mathrm{Tr}(XYZ) \leq ||XYZ||_1 \leq ||X||_p||Y||_q||Z||_r.$$

We also need the famous Lieb-Thirring inequality: $\mathrm{Tr}((AB)^m) \leq \mathrm{Tr}(A^m B^m)$.

Theorem 2.3. *Let X, Y be positive definite matrices and $z \in S_{1/4} = \{z \in \mathbb{C} : Re(z) \in [\frac{1}{4}, \frac{3}{4}]\}$. Then*

$$|\,\mathrm{Tr}(X^{1/2}Y^z Y^{1/2}Y^{1-z})| \leq \mathrm{Tr}(XY). \tag{12}$$

Proof. Let $z = \frac{1}{2} + iy, y \in \mathbb{R}$ denote any point in the vertical line of the complex plane passing $x = 1/2$. Then we have

$$\begin{aligned}|\,\mathrm{Tr}(X^{1/2}Y^z X^{1/2}Y^{1-z})| &= |\,\mathrm{Tr}(X^{1/2}Y^{1/2}Y^{iy}X^{1/2}Y^{1/2}Y^{-iy})| \\ &\leq \mathrm{Tr}(|X^{1/2}Y^{1/2}Y^{iy}X^{1/2}Y^{1/2}Y^{-iy}|) \\ &\leq ||X^{1/2}Y^{1/2}Y^{iy}||_2||X^{1/2}Y^{1/2}Y^{-iy}||_2 \\ &= ||X^{1/2}Y^{1/2}||_2^2 \\ &= \mathrm{Tr}(XY).\end{aligned}$$

The first inequality is obvious, the second one follows from the Cauchy-Schwarz inequality for trace, and the second equality is from the fact that Y^{iy} and Y^{-iy} are unitary operators.

Now let consider $z = \frac{1}{4} + iy, y \in \mathbb{R}$, a generic point in the vertical line over $x = 1/4$, then by using the Hölder inequality with $\frac{1}{4} + \frac{1}{4} + \frac{1}{2} = 1$ and the Araki-Lieb-Thirring inequality we have

$$\begin{aligned}
|\operatorname{Tr}(X^{1/2}Y^zX^{1/2}Y^{1-z})| &= |\operatorname{Tr}(X^{1/2}Y^{1/4}Y^{iy}X^{1/2}Y^{1/2}Y^{-iy}Y^{1/4})| \\
&= |\operatorname{Tr}(Y^{1/4}X^{1/4}X^{1/4}Y^{1/4}Y^{iy}X^{1/2}Y^{1/2}Y^{-iy})| \\
&\leq ||Y^{1/4}X^{1/4}||_{1/4}^2||X^{1/2}Y^{1/2}||_2 \\
&\leq ||Y^{1/2}X^{1/2}||_2||X^{1/2}Y^{1/2}||_2 \\
&= \operatorname{Tr}(XY).
\end{aligned}$$

Mention that the map $x \mapsto A^z = e^{x\ln A} = \sum_k z^k \frac{(\ln A)^k}{k!}$ is analytic for $A > 0$, the product of matrices is also analytic and the trace is complex linear, the function $f(z) = \operatorname{Tr}(X^{1/2}Y^zX^{1/2}Y^{1-z})$ is entire. Moreover, by the similar above argument for $z = x + iy$ it is easy to see that if $0 \leq x \leq 1$ then the function is bounded. By the Hadamard three-lines theorem the supremum $M(x) = \sup\{|f(x + iy)| : y \in \mathbb{R}\}$ of the function $\operatorname{Tr}(X^{1/2}Y^zX^{1/2}Y^{1-z})$ is log-convex, that means, for any $\lambda \in [0, 1]$,

$$M(\lambda x_1 + (1-\lambda)x_2) \leq M(x_1)^\lambda M(x_2)^{1-\lambda} \leq \operatorname{Tr}(XY)^\lambda \operatorname{Tr}(XY)^{1-\lambda} = \operatorname{Tr}(XY).$$

Therefore, the bound $\operatorname{Tr}(XY)$ is valid in the vertical strip $1/4 \leq \operatorname{Re}(z) \leq 1/2$. Invoking the symmetry $z \mapsto 1 - z$ and exchanging the roles of A and B give the desired bound on the full strip $S_{1/4} = \{1/4 \leq \operatorname{Re}(z) \leq 3/4\}$. □

As a consequence, we have the following inequality (see [1, Inequality (39)]):

Corollary 2.4. *For any positive definite matrices and for $t \in [0, 1]$.*

$$\operatorname{Tr}((A\sharp_t B)(A\sharp_{1-t}B)) \leq \operatorname{Tr}(AB).$$

Now we are ready to prove the main result in this paper.

Theorem 2.5. *Let A, B be positive definite matrices, $p = 1, 2$ and $r \geq 0$. Then*

$$||A + B + r(A\sharp_t B + A\sharp_{1-t}B)||_p \leq ||A + B + r(A^tB^{1-t} + A^{1-t}B^t)||_p. \tag{13}$$

Proof. Since $A+B+r(A\sharp_t B+A\sharp_{1-t}B) \geq 0$, the left hand side of (13) is $\operatorname{Tr}(A+B+r(A\sharp_t B+A\sharp_{1-t}B))$. It is well-known that $\operatorname{Tr}(A\sharp_t B) \leq \operatorname{Tr}(A^{1-t}B^t)$ and $\operatorname{Tr}(A\sharp_{1-t}B) \leq \operatorname{Tr}(A^tB^{1-t})$. We have

$$\begin{aligned}
\operatorname{Tr}(A + B + r(A\sharp_t B + A\sharp_{1-t}B)) &\leq \operatorname{Tr}(A + B + r(A^tB^{1-t} + A^{1-t}B^t)) \\
&\leq \operatorname{Tr}(|A + B + r(A^tB^{1-t} + A^{1-t}B^t)|).
\end{aligned}$$

So for $p = 1$ the inequality (13) follows.

Next consider the case $p = 2$. Notice again that $\mathrm{Tr}((A\sharp_t B)^2) \leq \mathrm{Tr}(B^{2t}A^{2(1-t)})$ (see [1, pape 121]). Similarly, we also have $\mathrm{Tr}((A\sharp_{1-t}B)^2) \leq \mathrm{Tr}(A^{2t}B^{2(1-t)})$. Then

$$\mathrm{Tr}((A\sharp_t B)^2 + (A\sharp_{1-t}B)^2) \leq \mathrm{Tr}(A^{2t}B^{2(1-t)} + B^{2t}A^{2(1-t)}). \tag{14}$$

By Proposition 2.1 we have

$$\mathrm{Tr}((A+B)(A\sharp_t B + A\sharp_{1-t}B)) \leq \mathrm{Tr}(A^{t+1}B^{1-t} + A^{2-t}B^t + A^tB^{2-t} + A^{1-t}B^{1+t}). \tag{15}$$

Now, squaring both sides of (13), we need to show

$$\begin{aligned}&\mathrm{Tr}((A+B)^2 + r^2(A\sharp_t B)^2 + r^2(A\sharp_{1-t}B)^2 + 2r(A+B)(A\sharp_t B + A\sharp_{1-t}B) + 2r^2(A\sharp_t B)(A\sharp_{1-t}B))\\ &\leq \mathrm{Tr}((A+B)^2 + 2r(A^{t+1}B^{1-t} + A^{2-t}B^t + A^tB^{2-t} + A^{1-t}B^{1+t}) + r^2A^{2t}B^{2(1-t)}) + r^2B^{2t}A^{2(1-t)}\\ &\quad + 2r^2\,\mathrm{Tr}(AB).\end{aligned}$$

The last inequality follows from (14), (15) and Corollary 2.4. □

Remark 2.6. From Theorem 2.5 for $s \in [0,1]$ we

$$||(1-s)(A+B) + s(A\sharp_t B + A\sharp_{1-t}B)||_p \leq ||(1-s)(A+B) + s(A^tB^{1-t} + A^{1-t}B^t)||_p.$$

When $t = 1/2$ we obtain one kind of inequality for the matrix Heron mean

$$||\frac{1-s}{2}(A+B) + s(A\sharp B)||_p \leq ||\frac{1-s}{2}(A+B) + sA^{1/2}B^{1/2}||_p.$$

Remark 2.7. By the same arguments, one can show that

$$||A+B+A\sharp_t B + A\sharp_{1-t}B||_p \leq ||A+B+A^tB^{1-t} + A^{1-t}B^t||_p.$$

But is we use another version of the Heinz mean $(A^tB^{1-t} + B^tA^{1-t})/2$ and realize the same proof in Theorem 2.5 the inequality in Corollary 2.4 could be as follows

$$\mathrm{Tr}((A\sharp_t B)(B\sharp_t A)) \leq \mathrm{Re}\,\mathrm{Tr}(A^tB^tA^{1-t}B^{1-t}). \tag{16}$$

Notice that both sides are bounded by $\mathrm{Tr}(AB)$ but it is not clear that (16) is true or not.

From the proof of the main theorem, it is natural to ask the following question: Is it true that for $0 \leq X, Y \leq Z$ such that $X \prec_{\log} Y$

$$Z^{1/2}XZ^{1/2} \prec_{\log} Z^{1/2}YZ^{1/2}? \tag{17}$$

Unfortunately, the answer is negative. Indeed, let

$$X = \begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}, \quad Y = \begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}, \quad Z = \begin{pmatrix}1 & 0\\ 0 & 4\end{pmatrix}.$$

Now $s(X) = s(Y) = (1,1)$ but $s(Z^{1/2}XZ^{1/2}) = (1,2) \not\prec_w (\sqrt{2},\sqrt{2}) = s(Z^{1/2}YZ^{1/2})$. So it is not even true for diagonal positive definite matrices.

3. Determinant Inequality for the Heron mean

Let's recall a recent result of Audeanert [9]: for any positive semidefinite matrices A and B

$$\det(I + A\sharp B) \leq \det(I + A^{1/2}B^{1/2}). \tag{18}$$

The author used the well-known fact that $\lambda(A\sharp B) \prec_{\log} \lambda(A^{1/2}B^{1/2})$ and the function $\Phi(X) = \sum_{i=1}^{n} \log(1 + e^{x_i})$ $(X = (x_1, x_2, \cdots, x_n))$ is isotone (i.e. the function preserving weak majorization: $x \prec y \Rightarrow \Phi(x) \prec_w \Phi(y)$.)

In fact, for matrices A and B such that $\lambda(A) \prec_{\log} \lambda(B)$ we have

$$\det(I + A) \leq \det(I + B). \tag{19}$$

A useful characterization of isotone functions in the case $m = 1$ is as follows:

Lemma 3.1. *A differentiable function $\Phi : \mathbb{R}^n \to \mathbb{R}$ is isotope if and only if it satisfy*

(1) *Φ is permutation invariant;*

(2) *for all $X \in \mathbb{R}^n$ and for all i, j:*

$$(x_i - x_j)\big(\frac{\partial \Phi}{\partial x_i}(x) - \frac{\partial \Phi}{\partial x_j}(x)\big) \geq 0.$$

Do the similar argument as in [9] one can prove the following

$$\det(I + A\sharp_t B) \leq \det(I + A^{1-t}B^t). \tag{20}$$

Now we can use this fact to obtain some inequality for the Heron mean.

Theorem 3.2. *For any positive definite matrices A and B*

$$\det(P_t(A, B)) \leq \det(Q_t(A, B)). \tag{21}$$

Proof. The inequality (21) is equivalent to the following

$$\begin{aligned} \det{}^{1/t}(A^t + B^t) &= \det(A) \det{}^{1/t}(I + A^{-t/2}B^tA^{-t/2}) \\ &\geq \det(A\sharp_{1/t}(A + A\sharp_t B)) \\ &= \det(A) \cdot \det{}^{1/t}(I + (A^{-1/2}BA^{-1/2})^t) \end{aligned}$$

or

$$\det(I + A^{-t/2}B^tA^{-t/2}) \geq \det(I + (A^{-1/2}BA^{-1/2})^t). \tag{22}$$

By the Araki-Lieb-Thirring inequality we have

$$\lambda(I + (A^{-1/2}BA^{-1/2})^t) \prec_{\log} \lambda(I + A^{-t/2}B^tA^{-t/2}).$$

Therefore, the inequality (22) follows from the last inequality and (19). □

As a consequence, we obtain a determinant inequality for the Heron mean.

Corollary 3.3. *For any positive definite matrices A and B,*

$$\det(A+B+2(A\sharp B)) \leq \det(A+B+A^{1/2}B^{1/2}+A^{1/2}B^{1/2})).$$

References

[1] R.Bhatia, Y.Lim, T.Yamazaki. *Some norm inequalities for matrix means.* Linear Algebra Appl. **501** (2016) 112-122.

[2] Y.Lim, M.Pálfia, The matrix power means and the Karcher mean. J. Funct. Anal. **262** (2012) 1498-1514.

[3] Y.Lim, T.Yamazaki. *On some inequalities for the matrix power and Karcher means.* Linear Algebra Appl. **438** (2013) 325-346.

[4] T.Bottazzi, R.Elencwajg, G.Larotonda, A.Varela. *Inequalities related to Bourin and Heinz means with a complex parameter.* J. Math. Anal. Appl. **426** (2015) 765-773.

[5] B. Simon. *Trace Ideals and Their Applications.* Second edition, Mathematical Surveys and Monographs, vol. 120, American Mathematical Society, Providence, RI, 2005.

[6] R.Bhatia, *Interpolating the arithmetic-geometric mean inequality and its operator version.* LAA **413** (2006) 355-363.

[7] R. Bhatia. *Positive Definite Matrices.* Princeton University Press, New Jersey, 2007.

[8] K.Kubo, T.Ando. *Means of positive linear operators.* Math. Ann. **246** (1979/80) 205-224.

[9] K.M.R. Audenaert. *A Determinantal Inequality for the Geometric Mean with an Application in Diffusion Tensor Imaging.*

Division of Computational Mathematics and Engineering (CME), Institute for Computational Science (INCOS), Ton Duc Thang University, Ho Chi Minh City, Vietnam;, Faculty of Civil Engineering, Ton Duc Thang University, Vietnam.

E-mail address: dinhtrunghoa@tdt.edu.vn